

NCTS/Math
Technical Report
2019-005



OPTIMAL EMBEDDINGS INTO LORENTZ SPACES FOR SOME VECTOR DIFFERENTIAL OPERATORS VIA GAGLIARDO'S LEMMA

DANIEL SPECTOR AND JEAN VAN SCHAFTINGEN

ABSTRACT. We prove a family of Sobolev inequalities of the form

$$\|u\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n,V)} \leq \|A(D)u\|_{L^1(\mathbb{R}^n,E)}$$

where $A(D) : C_c^\infty(\mathbb{R}^n, V) \rightarrow C_c^\infty(\mathbb{R}^n, E)$ is a vector first-order homogeneous linear differential operator with constant coefficients, u is a vector field on \mathbb{R}^n and $L^{\frac{n}{n-1},1}(\mathbb{R}^n)$ is a Lorentz space. These new inequalities imply in particular the extension of the classical Gagliardo–Nirenberg inequality to Lorentz spaces originally due to Alvino and a sharpening of an inequality in terms of the deformation operator by Strauss (Korn–Sobolev inequality) on the Lorentz scale. The proof relies on a nonorthogonal application of the Loomis–Whitney inequality and Gagliardo's lemma.

1. INTRODUCTION AND MAIN RESULTS

A now classical result of Gagliardo [17] and Nirenberg [29] asserts the existence of a constant $C > 0$ such that the inequality

$$(1.1) \quad \|u\|_{L^{n/(n-1)}(\mathbb{R}^n)} \leq C \|Du\|_{L^1(\mathbb{R}^n, \mathbb{R}^n)}$$

holds for all $u \in W^{1,1}(\mathbb{R}^n)$. While optimal on the scale of Lebesgue spaces, one can improve the target to a better Lorentz space. Indeed, Alvino [2] proved that there exists a constant C' such that the inequality

$$(1.2) \quad \|u\|_{L^{n/(n-1),1}(\mathbb{R}^n)} \leq C' \|Du\|_{L^1(\mathbb{R}^n, \mathbb{R}^n)}$$

holds for all functions $u \in W^{1,1}(\mathbb{R}^n)$ (see Section 2 for a precise definition of the Lorentz space $L^{p,q}(\mathbb{R}^n)$), with an explicit optimal value of the constant C' . The estimate (1.2) reaches a limiting case of the class of Sobolev embeddings into Lorentz spaces treated by O'Neil and Peetre [30, §3; 31, Théorème 7.1] for $u \in W^{1,p}(\mathbb{R}^n)$, $p > 1$; the corresponding improved Lorentz estimates when $p = n$ lead to exponential integrability estimates [9, Theorem 7; 10, Theorems 2 and 3]. The inequality (1.2) was rediscovered by Poornima [32] and Tartar [34, Theorem 8], and was also proved by Fournier [16]. As $L^{n/(n-1),1}(\mathbb{R}^n) \subsetneq L^{n/(n-1),n/(n-1)}(\mathbb{R}^n) = L^{n/(n-1)}(\mathbb{R}^n)$, the inequality (1.2) improves

NATIONAL CHIAO TUNG UNIVERSITY, DEPARTMENT OF APPLIED MATHEMATICS, HSINCHU, TAIWAN
NATIONAL CENTER FOR THEORETICAL SCIENCES, NATIONAL TAIWAN UNIVERSITY, NO. 1 SEC. 4
ROOSEVELT RD., TAIPEI, 106, TAIWAN

WASHINGTON UNIVERSITY IN ST. LOUIS, DEPARTMENT OF MATHEMATICS, ONE BROOKINGS DRIVE,
ST. LOUIS, MO 63130-4899

UNIVERSITÉ CATHOLIQUE DE LOUVAIN, INSTITUT DE RECHERCHE EN MATHÉMATIQUE ET PHYSIQUE,
CHEMIN DU CYCLOTRON 2 BTE L7.01.01, 1348 LOUVAIN-LA-NEUVE, BELGIUM

(1.1), while simple examples show that one cannot obtain further improvement in the second parameter. That one should be interested in the Lorentz spaces in general, or the sharpening of the inequality (1.1) found in (1.2) in particular, can be seen from a number of perspectives. A first motivation comes from real interpolation of Banach spaces [3, 18, 23–25], in which the Lorentz spaces arise readily

$$L^{p,q}(\mathbb{R}^n) = (L^1(\mathbb{R}^n), L^\infty(\mathbb{R}^n))_{1-\frac{1}{p}, q},$$

(see, e.g. [1, Theorem 7.26; 25, Remarque (2.1); 35, Lemma 22.6; 37, Théorème 1.18.16.1]), while the weak- L^p space $L^{p,\infty}(\mathbb{R}^n)$ is a natural space for many harmonic analysis estimates, where weak-type endpoints can be upgraded to strong-type interpolated estimates. In our specific considerations, the improvement in the second parameter is more than microscopic, as it encodes significantly more information in the trade off between differentiability and integrability than the classical inequality (1.1). One perspective of this gain is that from (1.2) it is possible to deduce Hardy's inequality

$$(1.3) \quad \int_{\mathbb{R}^n} \frac{|u(x)|}{|x|} dx \leq \tilde{C} \int_{\mathbb{R}^n} |Du(x)| dx,$$

by a simple application of Hölder's inequality on the Lorentz space scale.

A vector analogue of (1.1) follows easily from the same argument, yet such an inequality is not optimal, as one does not need the full gradient in order to obtain an embedding into $L^{n/(n-1)}(\mathbb{R}^n)$. For example, a result of M.J. Strauss [33] shows that if one defines the symmetric part of the gradient

$$Eu \triangleq \frac{1}{2}(Du + (Du)^T)$$

(Eu is known in elasticity as the linearized deformation tensor associated to the displacement u), then one has the existence of a constant $C'' > 0$ such that

$$(1.4) \quad \|u\|_{L^{n/(n-1)}(\mathbb{R}^n)} \leq C'' \|Eu\|_{L^1(\mathbb{R}^n, \mathbb{R}^n)}$$

for all vector fields $u \in W^{1,1}(\mathbb{R}^n, \mathbb{R}^n)$. The need for such inequalities arose in the work of Duvaut and Lions [13], while interest in the study of such spaces has expanded greatly into the theory of functions of bounded deformation [4, 5, 11, 36]. The inequality (1.4) is a special application of Strauss's work, which can be deduced from a more refined inequality [33, p. 208] in the spirit of a preceding work in the L^2 case due to De Figueiredo [12].

Similar estimates have been proved for other differential operators, including the Hodge complex, in a series of works initiated by Bourgain and Brezis (see [6, 7, 22, 38]), while more generally, the second author has shown that the vector differential inequality

$$(1.5) \quad \|u\|_{L^{\frac{n}{n-1}}(\mathbb{R}^n, V)} \leq \tilde{C} \|A(D)u\|_{L^1(\mathbb{R}^n, E)}$$

holds for every vector field $u \in C_c^\infty(\mathbb{R}^n, V)$ if and only if the homogeneous first-order linear vector differential operator with constant coefficients $A(D)$ is elliptic and canceling [41, Theorem 1.3] (see also Section 5.3).

Given the sharpening of the inequality (1.1) obtained in (1.2), a natural question is whether all elliptic and canceling operators admit such improvements (see [7, Open problem 1; 40, Open problem 2; 41, Open problem 8.3; 42, Open problem 2]). By relying on estimates into fractional Sobolev spaces and embeddings of these into Lorentz spaces, it

has been proved that such improvements can be obtained in the Lorentz space $L^{\frac{n}{n-1},q}(\mathbb{R}^n)$ for every $q > 1$ [40, Theorem 3; 41, Theorem 8.5]. The possibility of extending (1.2) to elliptic and canceling operators is supported by the fact that the analogue of the Hardy inequality (1.3) for such operators that it would imply is known to hold [8].

The main result of this paper is the positive answer to such a result in the plane, and a partial answer in higher dimensions, that such an embedding holds for elliptic and $(n-1)$ -canceling operators. From this we show how one can deduce the inequality (1.2), as well as the following sharpening of Strauss' result (1.4).

Theorem 1. *There exists a constant $C > 0$ such that*

$$\|u\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n,\mathbb{R}^n)} \leq C \|Eu\|_{L^1(\mathbb{R}^n,\mathbb{R}^{n \times n})}$$

for every $u \in C_c^\infty(\mathbb{R}^n, \mathbb{R}^n)$.

Luc Tartar mentioned in 2012 to the second author that he had a proof of Theorem 1 that has not yet been published; it appeared afterwards that both our independent proofs of Theorem 1 were following the same strategy.

The idea underlying the improvement of Theorem 1 is contained in the following general theorem which allows one to control the Lorentz norm by a product of directional derivatives.

Theorem 2. *Let $\ell \in \mathbb{N}$, let, for $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, \ell\}$, $w_i^j \in \mathbb{R}^n$ and $v_i^j \in V^*$. If for every $j \in \{1, \dots, \ell\}$, the vectors w_1^j, \dots, w_n^j are linearly independent in \mathbb{R}^n and if*

$$\bigcap_{j=1}^{\ell} \bigcup_{i=1}^n (v_i^j)^\perp = \{0\},$$

then for every function $u \in C_c^\infty(\mathbb{R}^n, V)$,

$$\|u\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n,V)} \leq C \sum_{j=1}^{\ell} \|\langle v_1^j, Du[w_1^j] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \cdots \|\langle v_n^j, Du[w_n^j] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}}.$$

From Theorem 2, we obtain a complete answer in the two-dimensional case:

Theorem 3. *Let V and E be finite-dimensional spaces and let $A(D) : C_c^\infty(\mathbb{R}^2, V) \rightarrow C_c^\infty(\mathbb{R}^2, E)$ be a first-order homogeneous linear differential operator with constant coefficients. There exists a constant $C > 0$ such that*

$$\|u\|_{L^{2,1}(\mathbb{R}^2,V)} \leq C \|A(D)u\|_{L^1(\mathbb{R}^2,E)}$$

for every $u \in C_c^\infty(\mathbb{R}^2, V)$ if and only if the operator $A(D)$ is elliptic and canceling.

Theorem 3 shows in particular that when $n = 2$ the Sobolev inequality (1.5) holds if and only if the corresponding limiting estimate in Lorentz spaces holds.

The proof of Theorem 3 in Section 5.3 shows in higher dimensions $n > 2$ that one has such inequalities for all elliptic and $(n-1)$ -canceling operators.

Theorem 3 motivates the following

Question 1.1. *Let $n \geq 3$ and let V and E be finite-dimensional spaces. Further suppose that the first-order homogeneous linear differential operator with constant coefficients $A(D) : C_c^\infty(\mathbb{R}^n, V) \rightarrow C_c^\infty(\mathbb{R}^n, E)$ is elliptic and canceling. Can one show the existence of a constant $C > 0$ such that the inequality*

$$\|u\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n, V)} \leq C \|A(D)u\|_{L^1(\mathbb{R}^n, E)}$$

holds for every $u \in C_c^\infty(\mathbb{R}^n, V)$?

Another open question stems from the fact that higher-order operators satisfy Sobolev estimates [41, Theorem 1.3], Hardy inequalities [8, Theorem 1.2] and non-optimal Lorentz estimates [41, Theorem 8.5].

Question 1.2. *Let $n \geq 2$, $k \geq 2$ and let V and E be finite-dimensional spaces. Further suppose that the k -th order homogeneous linear differential operator with constant coefficients $A(D) : C_c^\infty(\mathbb{R}^n, V) \rightarrow C_c^\infty(\mathbb{R}^n, E)$ is elliptic and canceling. Can one show the existence of a constant $C > 0$ such that the inequality*

$$\|D^{k-1}u\|_{L^{\frac{n}{n-1},1}(\mathbb{R}^n, V)} \leq C \|A(D)u\|_{L^1(\mathbb{R}^n, E)}$$

holds for every $u \in C_c^\infty(\mathbb{R}^n, V)$?

The plan of the paper is as follows. In Section 2, we recall some requisite preliminaries regarding Lorentz spaces. In Section 3, we give a proof of Theorem 1 in the plane, as it illustrates well the idea of the more general Theorem 2. The main ingredient for higher dimensions is a version of an inequality of Loomis and Whitney [26] to nonorthogonal coordinate systems. We show how this can be obtained from a change of coordinates and a Lemma of Gagliardo in [17] in a presentation that tries to keep the geometric content of the inequality. In Section 5.1 we prove Theorem 2, from which Theorem 1 is deduced in Section 5.2. Finally, in Section 5.3 we prove a general result for elliptic and $(n-1)$ -canceling operators.

2. PRELIMINARIES

In the sequel, with an abuse of notation we utilize $|\cdot|$ to denote the norm in any finite-dimensional vector space, e.g. the absolute value, the norm in Euclidean space \mathbb{R}^n , and the norm in V and E .

For $n \in \mathbb{N}$, $1 < p < +\infty$ and $1 \leq q \leq +\infty$, we denote by $L^{p,q}(\mathbb{R}^n)$ the Lorentz space [27] (see also for example [19]) with quasinorm

$$(2.1) \quad \|u\|_{L^{p,q}(\mathbb{R}^n)}^q \triangleq p \int_0^{+\infty} (t \mathcal{L}^n(\{x \in \mathbb{R}^n : |u(x)| > t\})^{1/p})^q \frac{dt}{t},$$

where $\mathcal{L}^n(A)$ denotes the Lebesgue measure of a measurable set $A \subset \mathbb{R}^n$. Equivalently, if the function $|u|^* : (0, +\infty) \rightarrow [0, +\infty)$ is the nonincreasing rearrangement of $|u|$, that is, if for every $t \in (0, +\infty)$, $\mathcal{L}^1(\{s \in [0, +\infty) : |u|^*(s) > t\}) = \mathcal{L}^n(\{x \in \mathbb{R}^n : |u(x)| > t\})$, then

$$\|u\|_{L^{p,q}(\mathbb{R}^n)}^q = p \int_0^{+\infty} (t \mathcal{L}^1(\{s \in (0, +\infty) : |u|^*(s) > t\})^{1/p})^q \frac{dt}{t} = \int_0^{+\infty} |u|^*(s) s^{\frac{q}{p}} \frac{ds}{s}.$$

In particular, Cavalieri's principle shows that $L^{p,p}(\mathbb{R}^n) = L^p(\mathbb{R}^n)$, while the spaces are nested increasingly with respect to the second parameter:

$$L^{p,1}(\mathbb{R}^n) \subset L^{p,q}(\mathbb{R}^n) \subset L^{p,\infty}(\mathbb{R}^n).$$

The quantity $\|\cdot\|_{L^{p,q}(\mathbb{R}^n)}$ is a norm when $q \leq p$ [27, Theorem 1].

The definition of Lorentz spaces and norms by (2.1) is equivalent when $p > 1$ and $q \geq 1$ to the definition by interpolation (see for example [1, Theorem 7.26; 25, Remarque (2.1); 35, Lemma 22.6; 37, Théorème 1.18.16.1]) and to the definition through *averaged rearrangements* which is also common [43, §1.8].

3. PROOF OF THEOREM 1 IN THE PLANAR CASE

We here give a proof of Theorem 1 in the plane \mathbb{R}^2 . That is, we suppose that the function $u : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is smooth and has compact support, and we will obtain an estimate for

$$\|u_1\|_{L^{2,1}(\mathbb{R}^2)} = 2 \int_0^\infty \mathcal{L}^2(\{x \in \mathbb{R}^2 : |u_1(x)| > t\})^{\frac{1}{2}} dt$$

by the quantity

$$\int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1} \right| + \left| \frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right| + \left| \frac{\partial u_2}{\partial x_2} \right|,$$

as an analogous argument implies a similar inequality for u_2 .

First let us recall Fournier's argument [16, Appendix] of how to use the Loomis–Whitney inequality to obtain the embedding (1.2) where one assumes the full derivative Du is in L^1 . In our setting of the plane this reduces to the degenerate case of the Loomis–Whitney inequality that the area A of a set can be bounded by the product of its length l and width w :

$$(3.1) \quad A \leq l \times w.$$

More specific to our problem, this takes for every $t > 0$ the form of the inequality

$$\begin{aligned} & \mathcal{L}^2(\{|u_1| > t\}) \\ & \leq \mathcal{H}^1\left(\left\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t\right\}\right) \mathcal{H}^1\left(\left\{x_2 \in \mathbb{R} : \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| > t\right\}\right), \end{aligned}$$

where \mathcal{H}^1 denotes the one-dimensional Hausdorff measure and $\{|u_1| > t\} = \{(x_1, x_2) \in \mathbb{R}^2 : |u_1(x_1, x_2)| > t\}$. One then observes that a separate integration of the two terms on the right-hand side yields

$$\int_0^\infty \mathcal{H}^1\left(\left\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t\right\}\right) dt = \int_{\mathbb{R}} \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| dx_1$$

and

$$\int_0^\infty \mathcal{H}^1\left(\left\{x_2 \in \mathbb{R} : \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| > t\right\}\right) dt = \int_{\mathbb{R}} \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| dx_2,$$

which can be further estimated by the derivatives via the fundamental theorem of calculus as

$$(3.2) \quad \int_{\mathbb{R}} \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| dx_1 \leq \int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_2}(x_1, x_2) \right| dx_2 dx_1,$$

and

$$(3.3) \quad \int_{\mathbb{R}} \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| dx_2 \leq \int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1}(x_1, x_2) \right| dx_1 dx_2.$$

Therefore, from the definition of the Lorentz norm (2.1) one has

$$\begin{aligned} \int_0^\infty \mathcal{L}^2(\{|u_1| > t\})^{\frac{1}{2}} dt &\leq \int_0^\infty \left(\mathcal{H}^1(\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t\}) \right. \\ &\quad \left. \times \mathcal{H}^1(\{x_2 \in \mathbb{R} : \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| > t\}) \right)^{\frac{1}{2}} dt, \end{aligned}$$

while the Cauchy–Schwarz inequality implies

$$(3.4) \quad \int_0^\infty \mathcal{L}^2(\{|u_1| > t\})^{\frac{1}{2}} dt \leq \left(\int_0^\infty \mathcal{H}^1(\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t\}) dt \right)^{\frac{1}{2}} \\ \times \left(\int_0^\infty \mathcal{H}^1(\{x_2 \in \mathbb{R} : \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| > t\}) dt \right)^{\frac{1}{2}}.$$

Finally one combines the inequalities (3.2), (3.3) and (3.4) to deduce the multiplicative inequality

$$\int_0^\infty \mathcal{L}^2(\{|u_1| > t\})^{\frac{1}{2}} dt \leq \left(\int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_2}(x_1, x_2) \right| dx_2 dx_1 \times \int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1}(x_1, x_2) \right| dx_1 dx_2 \right)^{\frac{1}{2}},$$

while the arithmetic geometric mean inequality yields the additive form

$$\int_0^\infty \mathcal{L}^2(\{|u_1| > t\})^{\frac{1}{2}} dt \leq \frac{1}{2} \int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_2}(x_1, x_2) \right| dx_2 dx_1 + \frac{1}{2} \int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1}(x_1, x_2) \right| dx_1 dx_2.$$

Now, this argument is not sufficient to obtain the Korn–Sobolev inequality, since in general one has no control over the quantity

$$\int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_2}(x_1, x_2) \right| dx_2 dx_1.$$

However, in this setting one still assumes the finiteness of

$$\int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1}(x_1, x_2) \right| dx_2 dx_1,$$

which tracing back through the inequalities translates to control over the width w in (3.1). In general we cannot hope to control the length l in this way, but it turns out we can control measurements in certain other directions. In particular, we can estimate the measurement of length in both directions whose angle with the x_2 axis is $\pi/4$. In

either case the measurement of l' gives us an upper bound on an estimate for l by simple trigonometry, leading to the inequality

$$(3.5) \quad A \leq \sqrt{2} l' \times w,$$

which as we will see will be sufficient to obtain our result.

We now commence with the

Proof of Theorem 1 in the planar case. Let us now see how this ability to control the area with respect to nonorthogonal measurements yields the desired inequality. First, we note that since, by the triangle inequality,

$$(3.6) \quad |u_1| \leq \frac{|u_1 + u_2| + |u_1 - u_2|}{2},$$

on \mathbb{R}^2 , we have for every $t > 0$

$$\{|u_1| > t\} \subset \{|u_1| > t, |u_1 + u_2| > t\} \cup \{|u_1| > t, |u_1 - u_2| > t\},$$

and therefore subadditivity of the measure \mathcal{L}^2 and of the square root implies that

$$(3.7) \quad \mathcal{L}^2(\{|u_1| > t\})^{\frac{1}{2}} \leq \mathcal{L}^2(\{|u_1| > t, |u_1 + u_2| > t\})^{\frac{1}{2}} + \mathcal{L}^2(\{|u_1| > t, |u_1 - u_2| > t\})^{\frac{1}{2}}.$$

Let us estimate the first term on the right-hand side. We apply the inequality (3.5) to deduce that for each $t > 0$,

$$\begin{aligned} & \mathcal{L}^2(\{|u_1| > t, |u_1 + u_2| > t\}) \\ & \leq \sqrt{2} \mathcal{H}^1\left(\left\{s \in \mathbb{R} : \sup_{x_1 - x_2 = s} |u_1| > t, \sup_{x_1 - x_2 = s} |u_1 + u_2| > t\right\}\right) \\ & \quad \times \mathcal{H}^1\left(\left\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1| > t, \sup_{x_2 \in \mathbb{R}} |u_1 + u_2| > t\right\}\right). \end{aligned}$$

Then the removal of certain inequalities in the sets only increases the measure, we find for each $t > 0$

$$\begin{aligned} \mathcal{L}^2(\{|u_1| > t, |u_1 + u_2| > t\}) & \leq \sqrt{2} \mathcal{H}^1\left(\left\{s \in \mathbb{R} : \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| > t\right\}\right) \\ & \quad \times \mathcal{H}^1\left(\left\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t, \right\}\right). \end{aligned}$$

Now while the integral in t of the second term on the right has been computed, for the first we find

$$\begin{aligned} & \int_0^\infty \mathcal{H}^1\left(\left\{s \in \mathbb{R} : \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| > t\right\}\right) dt \\ & = \int_{\mathbb{R}} \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| ds. \end{aligned}$$

We claim that this diagonal length can be controlled by the symmetric part of the gradient via the estimate

$$(3.8) \quad \int_{\mathbb{R}} \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| ds \leq \int_{\mathbb{R}^2} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} + \frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right) (x_1, x_2) \right| dx_1 dx_2,$$

from which the desired bound can be deduced, as the Cauchy–Schwarz inequality yields

$$\begin{aligned} & \int_0^\infty \mathcal{L}^2(\{|u_1| > t, |u_1 + u_2| > t\})^{\frac{1}{2}} dt \\ & \leq \left(\int_0^\infty \mathcal{H}^1\left(\left\{x_1 \in \mathbb{R} : \sup_{x_2 \in \mathbb{R}} |u_1(x_1, x_2)| > t\right\}\right) dt \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_0^\infty \mathcal{H}^1\left(\left\{s \in \mathbb{R} : \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| > t\right\}\right) dt \right)^{\frac{1}{2}}, \end{aligned}$$

and therefore

$$\begin{aligned} & \int_0^\infty \mathcal{L}^2(\{|u_1| > t, |u_1 + u_2| > t\})^{\frac{1}{2}} dt \\ & \leq \left(\int_{\mathbb{R}} \sup_{x_1 \in \mathbb{R}} |u_1(x_1, x_2)| dx_1 \right)^{\frac{1}{2}} \times \left(\int_{\mathbb{R}} \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| ds \right)^{\frac{1}{2}} \\ & \leq \left(\int_{\mathbb{R}^2} \left| \frac{\partial u_1}{\partial x_1}(x_1, x_2) \right| dx_2 dx_1 \right)^{\frac{1}{2}} \\ & \quad \times \left(\int_{\mathbb{R}^2} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} + \frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right)(x_1, x_2) \right| dx_1 dx_2 \right)^{\frac{1}{2}}. \end{aligned}$$

It therefore remains to prove the claim (3.8), as well as a similar estimate relating to a bound for the measure of the set $\{|u_1| > t, |u_1 - u_2| > t\}$. These two estimates are achieved by a modification of the argument of Gagliardo [17] and Nirenberg [29], that one can integrate in any direction and pair the gradient with an arbitrary covector (and we continue to restrict our consideration to the plane): For any vector $v \in \mathbb{R}^2 = (\mathbb{R}^2)^*$ and any vector $w \in \mathbb{R}^2$ one has

$$|\langle v, u(x_1, x_2) \rangle| \leq \int_{\mathbb{R}} |\langle v, Du(x + tw)w \rangle| dt.$$

Here again the choices $w = v = (1, 1)$ and $w = v = (1, -1)$ lead to the inequalities

$$|(u_1 + u_2)(x_1, x_2)| \leq \int_{\mathbb{R}} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} + \frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right)(x_1 + t, x_2 + t) \right| dt,$$

and

$$|(u_1 - u_2)(x_1, x_2)| \leq \int_{\mathbb{R}} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} - \left(\frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right) \right)(x_1 + t, x_2 - t) \right| dt.$$

Making a translation in t , one observes that the integrals on the right-hand-side depend only on $x_1 - x_2$ and $x_1 + x_2$, respectively. Letting $x_1 - x_2 = s \in \mathbb{R}$ in the former and $x_1 + x_2 = s \in \mathbb{R}$ in the latter, for each such s we can take the supremum over all such pairs (x_1, x_2) and then integrate in s to obtain

$$\int_{\mathbb{R}} \sup_{x_1 - x_2 = s} |(u_1 + u_2)(x_1, x_2)| ds \leq \int_{\mathbb{R}^2} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} + \frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right)(t, t - s) \right| dt ds$$

and

$$\int_{\mathbb{R}} \sup_{x_1+x_2=s} |(u_1 - u_2)(x_1, x_2)| ds \leq \int_{\mathbb{R}^2} \left| \left(\frac{\partial u_1}{\partial x_1} + \frac{\partial u_2}{\partial x_2} - \left(\frac{\partial u_1}{\partial x_2} + \frac{\partial u_2}{\partial x_1} \right) \right) (t, s - t) \right| dt ds.$$

It only remains to change variables to see that the claim has been demonstrated. \square

4. GAGLIARDO'S LEMMA AND LOOMIS–WHITNEY INEQUALITY

Our proof is based on a geometric inequality between the measure of a set and the measure of its projections on hyperplanes which goes back to Loomis and Whitney [26]. To state the following generalization of their inequality, we require the notion of $(n - 1)$ -dimensional Hausdorff measure of a set A , which we denote by $\mathcal{H}^{n-1}(A)$, see e.g. [14, §2.1, p. 60].

Lemma 4.1 (Loomis–Whitney inequality). *Let w_1, \dots, w_n be a basis of unit-length vectors of \mathbb{R}^n and let, for every $j \in \{1, \dots, n\}$, $\Pi_j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ denote the orthogonal projection of \mathbb{R}^n on w_j^\perp . Then there exists a constant such that for every compact set $K \subset \mathbb{R}^n$,*

$$\mathcal{L}^n(K)^{n-1} \leq \frac{\mathcal{H}^{n-1}(\Pi_1(K)) \cdots \mathcal{H}^{n-1}(\Pi_n(K))}{|\det(w_1, \dots, w_n)|}.$$

In the two-dimensional plane, the constant appearing in the inequality corresponds geometrically to the absolute value of the sine of the angle between the vectors w_1 and w_2 , while in any number of dimensions we have that equality is achieved in Lemma 4.1 when K is a parallelepiped spanned by the vectors w_1, \dots, w_n .

The original statement of Loomis and Whitney assumes that the vectors w_1, \dots, w_n are the canonical basis of \mathbb{R}^n Lemma 4.1 and is proved by a combinatorial argument through an approximation by sets that are a finite collection of cubes. Our approach shows how not only can one obtain the result of Loomis and Whitney as a direct consequence of the particular case of characteristic functions of a later lemma of Gagliardo [17, lemma 4.1], in fact one easily obtains in a geometric fashion the preceding more general version of their result.

Lemma 4.2. *Let $n \geq 2$. Let P_i denote the canonical projection of \mathbb{R}^n on $\mathbb{R}^{n-1} \simeq \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i} \subset \mathbb{R}^n$. For every choice of $f_i \in L^{n-1}(\mathbb{R}^{n-1})$, $i \in \{1, \dots, n\}$, one has $\prod_{i=1}^n f_i \circ P_i \in L^1(\mathbb{R}^n)$ with the estimate*

$$\left| \int_{\mathbb{R}^n} \prod_{i=1}^n f_i \circ P_i \right| \leq \prod_{i=1}^n \left(\int_{\mathbb{R}^{n-1}} |f_i|^{n-1} \right)^{\frac{1}{n-1}}.$$

The proof is the classical proof of Gagliardo that we give here for the convenience of the reader.

Proof of Lemma 4.2. We proceed by induction. First let us treat the base case, $n = 2$. In this case, from an application of Fubini's theorem we find

$$\begin{aligned} \left| \int_{\mathbb{R}^2} (f_1 \circ P_1)(f_2 \circ P_2) \right| &\leq \int_{\mathbb{R}^2} |f_1(z_2)| |f_2(z_1)| dz_1 dz_2 \\ &= \left(\int_{\mathbb{R}} |f_1| \right) \left(\int_{\mathbb{R}} |f_2| \right). \end{aligned}$$

Thus we proceed to the general case. For $n \geq 3$, we assume the lemma has been proved for $n - 1$ and will prove it for n . By Fubini's theorem we have

$$\int_{\mathbb{R}^n} \prod_{i=1}^n f_i \circ P_i = \int_{\mathbb{R}^{n-1}} \left(\int_{\mathbb{R}} \prod_{i=1}^{n-1} f_i(P_i(z)) \, dz_n \right) f_n(z') \, dz'.$$

From two applications of Hölder's inequality successively on \mathbb{R}^{n-1} and on \mathbb{R} we deduce

$$\begin{aligned} & \left| \int_{\mathbb{R}^n} \prod_{i=1}^n f_i \circ P_i \right| \\ (4.1) \quad & \leq \left(\int_{\mathbb{R}^{n-1}} \left| \int_{\mathbb{R}} \prod_{i=1}^{n-1} f_i(P_i(z', z_n)) \, dz_n \right|^{\frac{n-1}{n-2}} \, dz' \right)^{\frac{n-2}{n-1}} \left(\int_{\mathbb{R}^{n-1}} |f_n|^{n-1} \right)^{\frac{1}{n-1}} \\ & \leq \left(\int_{\mathbb{R}^{n-1}} \left(\prod_{i=1}^{n-1} \int_{\mathbb{R}} |f_i(P_i(z', z_n))|^{n-1} \, dz_n \right)^{\frac{1}{n-2}} \, dz' \right)^{\frac{n-2}{n-1}} \left(\int_{\mathbb{R}^{n-1}} |f_n|^{n-1} \right)^{\frac{1}{n-1}}. \end{aligned}$$

We now work to apply our induction assumption. For each $i \in \{1, \dots, n-1\}$, we define the function $g_i : \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-1-i} \times \{0\} \simeq \mathbb{R}^{n-2} \rightarrow \mathbb{R}$ for each $y \in \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-1-i} \times \{0\}$ by

$$(4.2) \quad g_i(y) \triangleq \left(\int_{\mathbb{R}} |f_i(y + (0, \dots, 0, z_n))|^{n-1} \, dz_n \right)^{\frac{1}{n-2}},$$

so that for every $z' \in \mathbb{R}^{n-1} \simeq \mathbb{R}^{n-1} \times \{0\} \subset \mathbb{R}^n$,

$$g_i(P_i(z')) = \left(\int_{\mathbb{R}} |f_i(P_i(z', z_n))|^{n-1} \, dz_n \right)^{\frac{1}{n-2}}.$$

We observe that by Fubini's theorem

$$\begin{aligned} (4.3) \quad \int_{\mathbb{R}^{n-2}} g_i^{n-2} &= \int_{\mathbb{R}^{n-2}} \int_{\mathbb{R}} |f_i(y + (0, \dots, z_n))|^{n-1} \, dz_n \, dy \\ &= \int_{\mathbb{R}^{n-1}} |f_i|^{n-1} < +\infty, \end{aligned}$$

so that $g_i \in L^{n-2}(\mathbb{R}^{n-2})$. Therefore we may apply our induction assumption to deduce

$$(4.4) \quad \int_{\mathbb{R}^{n-1}} \prod_{i=1}^{n-1} (g_i \circ P_i) \leq \prod_{i=1}^{n-1} \left(\int_{\mathbb{R}^{n-2}} g_i^{n-2} \right)^{\frac{1}{n-2}}.$$

Putting these inequalities (4.1), (4.2) and (4.4) together we find

$$\begin{aligned}
& \left| \int_{\mathbb{R}^n} \prod_{i=1}^n f_i \circ P_i \right| \\
& \leq \left(\int_{\mathbb{R}^{n-1}} \prod_{i=1}^{n-1} \left(\int_{\mathbb{R}} |f_i(P_i(z', z_n))|^{n-1} dz_n \right)^{\frac{1}{n-2}} dz' \right)^{\frac{n-2}{n-1}} \left(\int_{\mathbb{R}^{n-1}} |f_n|^{n-1} \right)^{\frac{1}{n-1}} \\
& = \left(\int_{\mathbb{R}^{n-1}} \prod_{i=1}^{n-1} g_i \circ P_i \right)^{\frac{n-2}{n-1}} \left(\int_{\mathbb{R}^{n-1}} |f_n|^{n-1} \right)^{\frac{1}{n-1}} \\
& \leq \prod_{i=1}^{n-1} \left(\int_{\mathbb{R}^{n-2}} g_i^{n-2} \right)^{\frac{1}{n-1}} \left(\int_{\mathbb{R}^{n-1}} |f_n|^{n-1} \right)^{\frac{1}{n-1}} \\
& = \prod_{i=1}^n \left(\int_{\mathbb{R}^{n-1}} |f_i|^{n-1} \right)^{\frac{1}{n-1}},
\end{aligned}$$

in view of the identity (4.3), which is the thesis. \square

Proof of Lemma 4.1. Let $\{w_1, \dots, w_n\}$ be a basis of \mathbb{R}^n . We define for every $i \in \{1, \dots, n\}$ the function $f_i : \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i} \rightarrow \mathbb{R}$ for each $z' \in \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i}$ by

$$f_i(z') \triangleq \sup_{t \in \mathbb{R}} \chi_K \left(\sum_{j \neq i} z_j w_j + t w_i \right),$$

where $\chi_K : \mathbb{R}^n \rightarrow \mathbb{R}$ is the characteristic function of the set K . Then we observe that for any $z \in \mathbb{R}^n$ we have

$$\chi_K \left(\sum_{j=1}^n z_j w_j \right) \leq f_i(P_i(z)),$$

and, as both sides assume only the values 0 and 1, we have

$$\chi_K \left(\sum_{j=1}^n z_j w_j \right) \leq \prod_{i=1}^n f_i(P_i(z)).$$

It follows thus that

$$\begin{aligned}
\mathcal{L}^n(K) &= |\det(w_1, \dots, w_n)| \int_{\mathbb{R}^n} \chi_K \left(\sum_{j=1}^n z_j w_j \right) dz \\
&\leq |\det(w_1, \dots, w_n)| \int_{\mathbb{R}^n} \prod_{i=1}^n f_i(P_i(z)) dz.
\end{aligned}$$

We observe now that by Gagliardo's inequality (Lemma 4.2) we have

$$\int_{\mathbb{R}^n} \prod_{i=1}^n f_i \circ P_i \leq \prod_{i=1}^n \left(\int_{\mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i}} |f_i|^{n-1} \right)^{\frac{1}{n-1}}.$$

But now for $z' \in \mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i}$ we have

$$f_i(z') = \sup_{t \in \mathbb{R}} \chi_K \left(\sum_{j \neq i} z_j w_j + t w_i \right) = (\chi_{\Pi_i(K)} \circ \Pi_i) \left(\sum_{j \neq i} z_j w_j \right),$$

while

$$\mathcal{H}^{n-1}(\Pi_i(K)) = J_i \int_{\mathbb{R}^i \times \{0\} \times \mathbb{R}^{n-i}} (\chi_{\Pi_i(K)} \circ \Pi_i) \left(\sum_{j \neq i} z_j w_j \right) dz',$$

where J_i is the (constant) Jacobian of the linear map $z' \in \mathbb{R}^i \times \{0\} \times \mathbb{R}^{n-i} \mapsto \Pi_i(\sum_{j \neq i} z_j w_j)$ (see [14, §3.2]). We now compute this Jacobian: if $i = 1$, we have since $|w_1| = 1$, by elementary manipulations of lines and columns of determinants

$$\begin{aligned} J_1^2 &= \det \begin{pmatrix} (w_2 \cdot w_2) - (w_1 \cdot w_2)(w_1 \cdot w_2) & \dots & (w_2 \cdot w_n) - (w_1 \cdot w_2)(w_1 \cdot w_n) \\ \vdots & \ddots & \vdots \\ (w_n \cdot w_2) - (w_1 \cdot w_n)(w_1 \cdot w_2) & \dots & (w_n \cdot w_n) - (w_1 \cdot w_n)(w_1 \cdot w_n) \end{pmatrix} \\ &= \det \begin{pmatrix} 1 & 0 & \dots & 0 \\ 0 & (w_2 \cdot w_2) - (w_1 \cdot w_2)(w_1 \cdot w_2) & \dots & (w_2 \cdot w_n) - (w_1 \cdot w_2)(w_1 \cdot w_n) \\ \vdots & \vdots & \ddots & \vdots \\ 0 & (w_n \cdot w_2) - (w_1 \cdot w_n)(w_1 \cdot w_2) & \dots & (w_n \cdot w_n) - (w_1 \cdot w_n)(w_1 \cdot w_n) \end{pmatrix} \\ &= \det \begin{pmatrix} 1 & (w_1 \cdot w_2) & \dots & (w_1 \cdot w_n) \\ 0 & (w_2 \cdot w_2) - (w_1 \cdot w_2)(w_1 \cdot w_2) & \dots & (w_2 \cdot w_n) - (w_1 \cdot w_2)(w_1 \cdot w_n) \\ \vdots & \vdots & \ddots & \vdots \\ 0 & (w_n \cdot w_2) - (w_1 \cdot w_n)(w_1 \cdot w_2) & \dots & (w_n \cdot w_n) - (w_1 \cdot w_n)(w_1 \cdot w_n) \end{pmatrix} \\ &= \det \begin{pmatrix} 1 & (w_1 \cdot w_2) & \dots & (w_1 \cdot w_n) \\ (w_2 \cdot w_1) & (w_2 \cdot w_2) & \dots & (w_2 \cdot w_n) \\ \vdots & \vdots & \ddots & \vdots \\ (w_n \cdot w_1) & (w_n \cdot w_2) & \dots & (w_n \cdot w_n) \end{pmatrix} \\ &= \det \begin{pmatrix} (w_1 \cdot w_1) & \dots & (w_1 \cdot w_n) \\ \vdots & \ddots & \vdots \\ (w_n \cdot w_1) & \dots & (w_n \cdot w_n) \end{pmatrix} = |\det(w_1, \dots, w_n)|^2 \end{aligned}$$

(this computation is in fact a case of computation of determinant through the Schur complement [21, (0.8.5.1)]); the case $i \in \{2, \dots, n\}$ is similar and therefore

$$\begin{aligned} \mathcal{L}^n(K) &\leq |\det(w_1, \dots, w_n)| \left(\prod_{i=1}^n \int_{\mathbb{R}^{i-1} \times \{0\} \times \mathbb{R}^{n-i}} \chi_{\Pi_i(K)} \left(\sum_{j \neq i} z_j w_j \right) dz' \right)^{\frac{1}{n-1}} \\ &= |\det(w_1, \dots, w_n)| \left(\prod_{i=1}^n \frac{\mathcal{H}^{n-1}(\Pi_i(K))}{|\det(w_1, \dots, w_n)|} \right)^{\frac{1}{n-1}} \\ &= \left(\frac{\prod_{i=1}^n \mathcal{H}^{n-1}(\Pi_i(K))}{|\det(w_1, \dots, w_n)|} \right)^{\frac{1}{n-1}}, \end{aligned}$$

thus concluding the demonstration of the claim. \square

5. PROOFS OF THE MAIN RESULTS

5.1. Estimates by directional derivatives of components. The last tool that we will need in the proofs is an estimate on the norm by sets of projections.

Lemma 5.1. *Under the assumptions of Theorem 2, there exists a constant $C \in \mathbb{R}$ such that for every $v \in V$, one has*

$$|v| \leq C \max \left\{ \min \{ |\langle v_1^j, v \rangle|, \dots, |\langle v_n^j, v \rangle| \} : j \in \{1, \dots, \ell\} \right\}.$$

Proof. Let $\gamma : V \rightarrow \mathbb{R}$ denote the function defined so that for every $v \in V$, the value $\gamma(v) \in \mathbb{R}$ is the right-hand side of the conclusion. The function γ is nonnegative, continuous and positively homogeneous of degree 1. We will reach the conclusion by proving that the function γ only vanishes at the point 0: indeed, if $v \in V$ and $\gamma(v) = 0$, then for every $j \in \{1, \dots, \ell\}$, we have $v \in \bigcup_{i=1}^n (v_i^j)^\perp$, and thus by assumption $v = 0$. \square

Proof of Theorem 2. For every $t > 0$, we have by Lemma 5.1,

$$\{x \in \mathbb{R}^n : |u(x)| \geq C_1 t\} \subseteq \bigcup_{j=1}^{\ell} \bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\},$$

for some constant $C_1 > 0$. We deduce then by subadditivity of the measure and of the map $\mu \in (0, +\infty) \mapsto \mu^{1-\frac{1}{n}}$ that

(5.1)

$$\begin{aligned} \mathcal{L}^n(\{x \in \mathbb{R}^n : |u(x)| \geq C_1 t\})^{1-\frac{1}{n}} &\leq \left(\sum_{j=1}^{\ell} \mathcal{L}^n \left(\bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\} \right) \right)^{1-\frac{1}{n}} \\ &\leq \sum_{j=1}^{\ell} \mathcal{L}^n \left(\bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\} \right)^{1-\frac{1}{n}}. \end{aligned}$$

If $j \in \{1, \dots, \ell\}$, then by assumption, the vectors w_1^j, \dots, w_n^j are linearly independent in \mathbb{R}^n and thus by the Loomis–Whitney inequality (Lemma 4.1) and by monotonicity of the measure, we have

$$\begin{aligned} &\mathcal{L}^n \left(\bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\} \right)^{1-\frac{1}{n}} \\ (5.2) \quad &\leq C_2 \prod_{k=1}^n \mathcal{H}^{n-1} \left(\Pi_k^j \left(\bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\} \right) \right)^{\frac{1}{n}} \\ &\leq C_2 \prod_{i=1}^n \mathcal{H}^{n-1} \left(\Pi_i^j (\{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\}) \right)^{\frac{1}{n}}, \end{aligned}$$

where $\Pi_i^j : \mathbb{R}^n \rightarrow \mathbb{R}^n$ is the orthogonal projection on the hyperplane $W_i^j \triangleq w_i^{j\perp} = \Pi_i^j(\mathbb{R}^n)$. Finally, we observe that if

$$y \in \Pi_i^j(\{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\}),$$

then there exists a real number $h \in \mathbb{R}$ such that $|\langle v_i^j, u(y + hw_i^j) \rangle| \geq t$ and thus

$$\int_{\mathbb{R}} |\langle v_i^j, Du(y + sw_i^j)[w_i^j] \rangle| ds \geq 2 |\langle v_i^j, u(y + hw_i^j) \rangle| \geq 2t.$$

If we define the function $F_i^j : W_i^j \rightarrow \mathbb{R}$ by setting for each $y \in W_i^j$

$$F_i^j(y) \triangleq \frac{1}{2} \int_{\mathbb{R}} |\langle v_i^j, Du(y + sw_i^j)[w_i^j] \rangle| ds,$$

we have for each $t > 0$

$$\Pi_i^j(\{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\}) \subseteq \{y \in W_i^j : F_i^j(y) \geq t\}$$

and thus by (5.2)

$$\mathcal{L}^n \left(\bigcap_{i=1}^n \{x \in \mathbb{R}^n : |\langle v_i^j, u(x) \rangle| \geq t\} \right)^{1-\frac{1}{n}} \leq C_2 \prod_{i=1}^n \left(\mathcal{H}^{n-1}(\{y \in W_i^j : F_i^j(y) \geq t\}) \right)^{\frac{1}{n}}.$$

In view of (5.1) and of the Hölder inequality, we obtain

$$\begin{aligned} & \int_0^\infty \mathcal{L}^n(\{x \in \mathbb{R}^n : |u(x)| \geq C_1 t\})^{1-\frac{1}{n}} dt \\ & \leq C_3 \sum_{j=1}^{\ell} \prod_{i=1}^n \left(\int_0^\infty \mathcal{H}^{n-1}(\{y \in W_i^j : F_i^j(y) \geq t\}) dt \right)^{\frac{1}{n}} \\ & = C_3 \sum_{j=1}^{\ell} \prod_{i=1}^n \left(\int_{W_i^j} F_i^j d\mathcal{H}^{n-1} \right)^{\frac{1}{n}} \\ & = \frac{C_3}{2} \sum_{j=1}^{\ell} \prod_{i=1}^n \left(\int_{\mathbb{R}^n} |\langle v_i^j, Du[w_i^j] \rangle| \right)^{\frac{1}{n}}. \quad \square \end{aligned}$$

5.2. Direct consequences. We firstly show of (1.2) can be deduced from Theorem 2.

Proof of (1.2) by Theorem 2. Let v_1, \dots, v_m be a basis of V^* and w_1, \dots, w_n be a basis of \mathbb{R}^n . We set $\ell = m$, and for every $i \in \{1, \dots, n\}$ and $j \in \{1, \dots, m\}$, $v_i^j = v_j$ and $w_i^j = w_i$. We have

$$\bigcap_{j=1}^m \bigcup_{i=1}^n (v_i^j)^\perp = \bigcap_{j=1}^m v_j^\perp = \{0\},$$

so that Theorem 2 applies and we conclude by Young's inequality and by norm equivalence that

$$\begin{aligned} \|u\|_{L^{\frac{n}{n-1}, 1}(\mathbb{R}^n, V)} & \leq C_1 \sum_{j=1}^m \prod_{i=1}^n \|\langle v_j, Du[w_i] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \\ & \leq \frac{C_1}{n} \sum_{j=1}^m \sum_{i=1}^n \|\langle v_j, Du[w_i] \rangle\|_{L^1(\mathbb{R}^n)} \leq C_2 \|Du\|_{L^1(\mathbb{R}^n, \mathbb{R}^n)}. \quad \square \end{aligned}$$

We next prove an analogue of De Figueiredo's L^2 inequality, from which we can deduce the improvement to Strauss' Korn-Sobolev inequality. To this end it will be useful to introduce the following definition.

Definition 5.2. If V is a vector space, then a finite set of vectors $F \subset V$ is *maximally linearly independent*, whenever for every subset $A \subset F$, either A generates V as a linear space or A is linearly independent.

Lemma 5.3. *If $n \geq 1$, $\dim V = m$ and if v_1, \dots, v_{n+m-1} are maximally linearly independent in V^* , then*

$$\bigcap_{1 \leq i_1 < \dots < i_n \leq n+m-1} \left(\bigcup_{j=1}^n v_{i_j}^\perp \right) = \{0\}.$$

Proof. Assume that v belongs to the left hand side and let $I = \{i \in \{1, \dots, n+m-1\} : \langle v_i, v \rangle = 0\}$. We have then $\#I \geq m$. Indeed, otherwise there would exist $i_1, \dots, i_n \in \{1, \dots, n+m-1\}$ such that $1 \leq i_1 < \dots < i_n \leq n+m-1$ and for every $j \in \{1, \dots, n\}$, $\langle v_{i_j}, v \rangle \neq 0$ and so $v \notin \bigcup_{j=1}^n v_{i_j}^\perp$, in contradiction with our assumption. Since the family v_1, \dots, v_{n+m-1} is maximally linearly independent, the set $\{v_i : i \in I\}$ generates the m -dimensional linear space V and thus $v = 0$. \square

Theorem 4. *Assume that $\dim V = m$ and that the vectors $w_1, \dots, w_{n+m-1} \in \mathbb{R}^n$ and $v_1, \dots, v_{n+m-1} \in V$ are maximally linearly independent, then for every $u \in C_c^\infty(\mathbb{R}^n, V)$,*

$$\|u\|_{L^{\frac{n}{n-1}, 1}(\mathbb{R}^n, V)} \leq \sum_{1 \leq i_1 < \dots < i_n \leq n+m-1} \|\langle v_{i_1}, Du[w_{i_1}] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \cdots \|\langle v_{i_n}, Du[w_{i_n}] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}}.$$

These sparse directional Sobolev estimates into Lorentz space are analogous to L^2 estimates of de Figueiredo [12] and strengthen known results for Sobolev estimates into $L^{\frac{n}{n-1}}$ [7, Remark 16; 33; 41, Proposition 6.8].

Proof of Theorem 4. This follows from Theorem 2 and Lemma 5.3. \square

Finally, we can utilize the preceding inequality to deduce the Korn–Sobolev inequality.

Proof of Theorem 1 by Theorem 4. We consider w_1, \dots, w_{2n-1} to be a maximally independent family of vectors of \mathbb{R}^n . We observe now that, since Eu is the symmetric part of Du ,

$$|\langle w_j, Du[w_j] \rangle| = |w_j \cdot Eu[w_j]| \leq |w_j|^2 |Eu|$$

and the conclusion then follows from Theorem 4. \square

5.3. Estimates for $(n-1)$ -canceling operators. In order to set an algebraic condition on differential operators, we introduce a new scale of conditions on differential operators that covers the definition of canceling operators [41, Definition 1.2].

Definition 5.4. Let $\ell \in \{0, \dots, n\}$. A homogeneous differential operator with constant coefficients $A(D)$ is ℓ -canceling whenever

$$\bigcap_{\substack{W \subseteq \mathbb{R}^n \\ \dim W = \ell}} \text{span} \{A(\xi)[v] : \xi \in W \text{ and } v \in V\} = \{0\}.$$

An operator is 1-canceling if and only if it is canceling in the sense of [41, Definition 1.2]. Any operator $A(D)$ is 0-canceling; an operator $A(D)$ is n -canceling if and only if $A(D) = 0$.

Optimal estimates into Lorentz spaces hold under the $(n-1)$ -canceling condition:

Theorem 5. *Let V and E be finite-dimensional spaces and suppose that the homogeneous linear differential operator with constant coefficients $A(D) : C_c^\infty(\mathbb{R}^n, V) \rightarrow C_c^\infty(\mathbb{R}^n, E)$ is elliptic and $(n-1)$ -canceling. Then there exists a constant $C > 0$ such that*

$$\|u\|_{L^{\frac{n}{n-1}, 1}(\mathbb{R}^n, V)} \leq C \|A(D)u\|_{L^1(\mathbb{R}^n, E)}$$

for every $u \in C_c^\infty(\mathbb{R}^n, V)$.

We will deduce Theorem 5 from Theorem 2 with the help of the next algebraic lemma.

Lemma 5.5. *If the first-order homogeneous differential operator with constant coefficients $A(D)$ is elliptic and $(n-1)$ -canceling, then there exists $m \in \mathbb{N}$ and, for each $j \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$, vectors $w_i^j \in \mathbb{R}^n$, $v_i^j \in V^*$ and $e_i^j \in E^*$ such that*

- (i) for every $j \in \{1, \dots, m\}$, the vectors w_1^j, \dots, w_n^j are linearly independent in \mathbb{R}^n ,
 - (ii) $\bigcap_{j=1}^m \bigcup_{i=1}^n (v_i^j)^\perp = \{0\}$,
 - (iii) for every $j \in \{1, \dots, m\}$, $i \in \{1, \dots, n\}$, $\xi \in \mathbb{R}^n$ and $v \in V$,
- $$\langle w_i^j, \xi \rangle \langle v_i^j, v \rangle = \langle e_i^j, A(\xi)[v] \rangle.$$

The proof of Lemma 5.5 will proceed by induction, the next lemma is the key step in the iteration.

Lemma 5.6. *If the first-order homogeneous differential operator with constant coefficients $A(D)$ is elliptic and canceling, then for every $v_* \in V \setminus \{0\}$, there exists vectors $\xi_1, \dots, \xi_n \in \mathbb{R}^n$, vectors $w_1, \dots, w_n \in \mathbb{R}^n$ and vectors $e_1, \dots, e_n \in E^*$ such that*

- (i) the vectors w_1, \dots, w_n are linearly independent in \mathbb{R}^n ,
- (ii) for every $i \in \{1, \dots, n\}$, $\langle e_i, A(\xi_i)[v_*] \rangle = 1$,
- (iii) for every $i \in \{1, \dots, n\}$, $\langle \xi_i, w_i \rangle = 1$,
- (iv) for every $i \in \{1, \dots, n\}$, $\xi \in \mathbb{R}^n$ and $v \in V$, $\langle e_i, A(\xi)[v] \rangle = \langle \xi, w_i \rangle \langle A(\xi_i)^*[e_i], v \rangle$.

Proof. We proceed by induction, that is, we are proving that for every $\ell \in \{0, 1, \dots, n\}$, there exists covectors $\xi_1, \dots, \xi_\ell \in \mathbb{R}^n$, vectors $w_1, \dots, w_\ell \in \mathbb{R}^n$ and vectors $e_1, \dots, e_\ell \in E^*$ such that

- (a) the vectors $w_1, \dots, w_\ell \in \mathbb{R}^n$ are linearly independent in \mathbb{R}^n ,
- (b) for every $i \in \{1, \dots, \ell\}$, $\langle e_i, A(\xi_i)[v_*] \rangle = 1$,
- (c) for every $i \in \{1, \dots, \ell\}$, $\langle \xi_i, w_i \rangle = 1$,
- (d) for every $i \in \{1, \dots, \ell\}$, $\xi \in \mathbb{R}^n$ and $v \in V$, $\langle e_i, A(\xi)[v] \rangle = \langle \xi, w_i \rangle \langle A(\xi_i)^*[e_i], v \rangle$.

For $\ell = 0$, the assertion holds vacuously. Assuming now that the assumption holds for some $\ell \in \{0, 1, \dots, n-1\}$, we will prove the assertion for $\ell + 1$.

Since $\ell \leq n-1$, there exists a covector $\xi_{\ell+1} \in \mathbb{R}^n \setminus \{0\}$ such that for every $i \in \{1, \dots, \ell\}$, $\langle \xi_{\ell+1}, w_i \rangle = 0$. (In particular, if $\ell = 0$, we just take any $\xi_1 \in \mathbb{R}^n \setminus \{0\}$.) Since the operator $A(D)$ is elliptic, the linear operator $A(\xi_{\ell+1}) : V \rightarrow E$ is injective, and thus, since $v_* \neq 0$, we have $A(\xi_{\ell+1})[v_*] \neq 0$. Since the operator $A(D)$ is also $(n-1)$ -canceling, there exists an $(n-1)$ -dimensional linear subspace $W_{\ell+1} \subset \mathbb{R}^n$ such that

$$(5.3) \quad A(\xi_{\ell+1})[v_*] \notin \text{span} \{A(\xi)[v] : \xi \in W_{\ell+1} \text{ and } v \in V\}.$$

We define now $w_{\ell+1} \in \mathbb{R}^n$ to be a vector such that $\langle \xi_{\ell+1}, w_{\ell+1} \rangle = 1$ and for every $\xi \in W_{\ell+1}$, one has $\langle \xi, w_{\ell+1} \rangle = 0$. In particular, this implies (c). Since the vectors w_1, \dots, w_ℓ

are linearly independent and since by construction we have for every $i \in \{1, \dots, \ell\}$, $\langle \xi_{\ell+1}, w_i \rangle = 0$, the vectors $w_1, \dots, w_\ell, w_{\ell+1}$ are linearly independent in \mathbb{R}^n and thus (a) holds. By (5.3) there exists a covector $e_{\ell+1} \in E^*$ such that $\langle e_{\ell+1}, A(\xi_{\ell+1})[v_*] \rangle = 1$ and for every $\xi \in W_{\ell+1}$ and $v \in V$, one has $\langle e_{\ell+1}, A(\xi)[v] \rangle = 0$. In particular, (b) holds. Moreover, we have $A(\xi)^*[e_{\ell+1}] = 0$ when $\xi \in W_{\ell+1}$. Therefore, since $\dim W_{\ell+1} = n - 1$, by the classical representation theorem of linear mappings, we deduce that for every $\xi \in \mathbb{R}^n$,

$$A(\xi)^*[e_{\ell+1}] = \langle \xi, w_{\ell+1} \rangle A(\xi_{\ell+1})^*[e_{\ell+1}],$$

which implies assertion (d). \square

A set $X \subseteq V$ is a *linear subspace arrangement* whenever X is a finite union of linear subspaces of V .

Lemma 5.7. *Let V be a finite-dimensional space. Assume that for each $\ell \in \mathbb{N}$, X_ℓ is a linear subspace arrangement of V and that $X_\ell \supseteq X_{\ell+1}$. Then there exists $\ell_0 \in \mathbb{N}$ such that for every $\ell \geq \ell_0$, $X_\ell = X_{\ell_0}$.*

Lemma 5.7 can be proved by observing that for every $\ell \in \mathbb{N}$, the set X_ℓ is Zariski-closed so that the sequence $(X_\ell)_{\ell \in \mathbb{N}}$ is a descending chain of Zariski-closed sets for which the conclusion follows (see for example [20, Example 1.4.7]). We give a direct proof for the convenience of the reader.

Proof of Lemma 5.7. For every $j \in \{1, \dots, \dim V\}$ we consider the set

$$X_\ell^j = \bigcup \{Y \subseteq V : Y \text{ is a linear subspace, } \dim Y \geq j \text{ and } Y \subseteq X_\ell\}.$$

We have $X_\ell^j \supseteq X_{\ell+1}^j$ and X_ℓ^j is a finite union of linear subspaces of dimension at least j .

We prove now by downward induction, that for every $j \in \{1, \dots, \dim V\}$, there exists ℓ_j such that for every $\ell \geq \ell_j$, we have $X_\ell^j = X_{\ell_j}^j$. For $j = \dim V$, either for every $\ell \in \mathbb{N}$, $X_\ell = V$ and then $\ell_j = 0$, or there exists $\ell_j \in \mathbb{N}$ such that $X_\ell = \{0\}$.

We assume now that the assertion is proved for some $j \in \{2, \dots, \dim V\}$. We observe that for every $\ell \geq \ell_j$, the components of X_ℓ^{j-1} of dimension at least j remain the same and the $(j-1)$ -dimensional components of X_ℓ^{j-1} form a subset of those of $X_{\ell_j}^{j-1}$. We have thus a nonincreasing sequence of finite subsets; there exists thus $\ell_{j-1} \geq \ell_j$ such that for every $\ell \geq \ell_{j-1}$, $X_\ell^{j-1} = X_{\ell_{j-1}}^{j-1}$. \square

We are now in position to prove Lemma 5.6.

Proof of Lemma 5.6. We are going to construct a the family of vectors iteratively over ℓ . At each step, we assume that we have vectors $w_i^j \in \mathbb{R}^n$, $v_i^j \in V^*$ and $e_i^j \in E^*$ for $j \in \{1, \dots, \ell\}$ and $i \in \{1, \dots, n\}$, such that

- (a) for every $j \in \{1, \dots, \ell\}$, the vectors w_1^j, \dots, w_n^j are linearly independent in \mathbb{R}^n ,
- (b) for every $j \in \{1, \dots, \ell\}$, $i \in \{1, \dots, n\}$, $\xi \in \mathbb{R}^n$ and $v \in V$,

$$\langle w_i^j, \xi \rangle \langle v_i^j, v \rangle = \langle e_i^j, A(\xi)[v] \rangle.$$

This is trivially satisfied when $\ell = 0$.

Assume thus that we have such families of vectors for some ℓ . If

$$X_\ell \triangleq \bigcap_{j=1}^{\ell} \bigcup_{i=1}^n (v_i^j)^\perp = \{0\},$$

then the proposition is proved with $m = \ell$. Otherwise, we take $v_*^{\ell+1} \in X_\ell \setminus \{0\}$, and we obtain by Lemma 5.6 vectors $\xi_1^{\ell+1}, \dots, \xi_n^{\ell+1} \in \mathbb{R}^n$, $w_1^{\ell+1}, \dots, w_n^{\ell+1} \in \mathbb{R}^n$ and $e_1^{\ell+1}, \dots, e_n^{\ell+1} \in E^*$. We set $v_i^{\ell+1} = A(\xi_i^{\ell+1})^*[e_i^{\ell+1}]$, and we observe that the family satisfies the same condition and moreover, since for every $i \in \{1, \dots, n\}$, $\langle v_i^{\ell+1}, v_*^{\ell+1} \rangle = \langle e_i^{\ell+1}, A(\xi_i^{\ell+1})[v_*^{\ell+1}] \rangle = 1$, we have $v_*^{\ell+1} \notin X_{\ell+1}$ and thus $X_{\ell+1} \subsetneq X_\ell$.

We conclude by observing that the procedure must finish after a finite number of steps in view of Lemma 5.7. \square

Proof of Theorem 5. Let $m \in \mathbb{N}$ and, for each $j \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$, the vectors $w_i^j \in \mathbb{R}^n$, $v_i^j \in V^*$ and $e_i^j \in E^*$ be given for $A(D)$ by Lemma 5.5. In view of Theorem 2, we have

$$\|u\|_{L^{\frac{n}{n-1}, 1}(\mathbb{R}^n, V)} \leq C \sum_{j=1}^m \|\langle v_1^j, Du[w_1^j] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}} \cdots \|\langle v_n^j, Du[w_n^j] \rangle\|_{L^1(\mathbb{R}^n)}^{\frac{1}{n}}.$$

Now the construction of Lemma 5.5 with $v = \widehat{u}(\xi)$, the Fourier transform of u at the point $\xi \in \mathbb{R}^n$, yields for every $j \in \{1, \dots, m\}$ and $i \in \{1, \dots, n\}$ and $\xi \in \mathbb{R}^n$,

$$\langle v_i^j, \langle 2\pi i \xi, w_i^j \rangle \widehat{u}(\xi) \rangle = \langle e_i^j, A(2\pi i \xi) \widehat{u}(\xi) \rangle,$$

and thus inverting the Fourier transform we obtain the pointwise equality

$$\langle v_i^j, Du[w_i^j] \rangle = \langle v_i^j, \langle Du, w_i^j \rangle \rangle = \langle e_i^j, A(D)u \rangle.$$

Hence

$$\|\langle v_i^j, Du[w_i^j] \rangle\|_{L^1(\mathbb{R}^n)} = \|\langle e_i^j, A(D)u \rangle\|_{L^1(\mathbb{R}^n)},$$

and the conclusion then follows. \square

Proof of Theorem 3. This follows from Theorem 5, [41, Theorem 1.3], the embedding between Lorentz spaces $L^{n/(n-1), 1}(\mathbb{R}^n) \subset L^{n/(n-1)}(\mathbb{R}^n)$ and the fact that the 1-canceling and canceling conditions are equivalent. \square

ACKNOWLEDGEMENTS

The authors would like to thank Wen-Wei Lin, the S.T. Yau Center at National Chiao Tung University, and the National Center for Theoretical Sciences of Taiwan for their support in the conference where this collaboration was initiated. The first author is supported in part by the Taiwan Ministry of Science and Technology under research grants 105-2115-M-009-004-MY2, 107-2918-I-009-003 and 107-2115-M-009-002-MY2.

REFERENCES

- [1] R. A. Adams and J. J. F. Fournier, *Sobolev spaces*, 2nd ed., Pure and Applied Mathematics (Amsterdam), vol. 140, Elsevier/Academic Press, Amsterdam, 2003. ↑2, 5
- [2] A. Alvino, *Sulla diseguglianza di Sobolev in spazi di Lorentz*, Boll. Un. Mat. Ital. A (5) **14** (1977), no. 1, 148–156. ↑1
- [3] N. Aronszajn and E. Gagliardo, *Interpolation spaces and interpolation methods*, Ann. Mat. Pura Appl. (4) **68** (1965), 51–117. ↑2
- [4] J.-F. Babadjian, *Traces of functions of bounded deformation*, Indiana Univ. Math. J. **64** (2015), no. 4, 1271–1290, doi:10.1512/iumj.2015.64.5601. ↑2
- [5] A. C. Barroso, I. Fonseca, and R. Toader, *A relaxation theorem in the space of functions of bounded deformation*, Ann. Scuola Norm. Sup. Pisa Cl. Sci. (4) **29** (2000), no. 1, 19–49. ↑2
- [6] J. Bourgain and H. Brezis, *New estimates for the Laplacian, the div-curl, and related Hodge systems*, C. R. Math. Acad. Sci. Paris **338** (2004), no. 7, 539–543, doi:10.1016/j.crma.2003.12.031. ↑2
- [7] ———, *New estimates for elliptic equations and Hodge type systems*, J. Eur. Math. Soc. (JEMS) **9** (2007), no. 2, 277–315, doi:10.4171/JEMS/80. ↑2, 15
- [8] P. Bousquet and J. Van Schaftingen, *Hardy–Sobolev inequalities for vector fields and canceling linear differential operators*, Indiana Univ. Math. J. **63** (2014), no. 5, 1419–1445, doi:10.1512/iumj.2014.63.5395. ↑3, 4
- [9] H. Brezis, *Laser beams and limiting cases of Sobolev inequalities*, Nonlinear partial differential equations and their applications. Collège de France Seminar, Vol. II (Paris, 1979), Res. Notes in Math., vol. 60, Pitman, Boston, Mass.-London, 1982, pp. 86–97. ↑1
- [10] H. Brezis and S. Wainger, *A note on limiting cases of Sobolev embeddings and convolution inequalities*, Comm. Partial Differential Equations **5** (1980), no. 7, 773–789. ↑1
- [11] G. Dal Maso, *Generalised functions of bounded deformation*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 5, 1943–1997, doi:10.4171/JEMS/410. ↑2
- [12] D. G. de Figueiredo, *The coerciveness problem for forms over vector valued functions*, Comm. Pure Appl. Math. **16** (1963), 63–94, doi:10.1002/cpa.3160160109. ↑2, 15
- [13] G. Duvaut and J.-L. Lions, *Un problème d'élasticité avec frottement*, J. Mécanique **10** (1971), 409–420. ↑2
- [14] L. C. Evans and R. F. Gariepy, *Measure theory and fine properties of functions*, Studies in Advanced Mathematics, CRC Press, Boca Raton, Fla., 1992. ↑9, 12
- [15] H. Federer and W. H. Fleming, *Normal and integral currents*, Ann. of Math. (2) **72** (1960), 458–520, doi:10.2307/1970227. ↑
- [16] J. J. F. Fournier, *Mixed norms and rearrangements: Sobolev's inequality and Littlewood's inequality*, Ann. Mat. Pura Appl. (4) **148** (1987), 51–76, doi:10.1007/BF01774283. ↑1, 5
- [17] E. Gagliardo, *Proprietà di alcune classi di funzioni in più variabili*, Ricerche Mat. **7** (1958), 102–137. ↑1, 4, 8, 9
- [18] ———, *Interpolazione di spazi di Banach e applicazioni*, Ricerche Mat. **9** (1960), 58–81. ↑2
- [19] L. Grafakos, *Classical Fourier analysis*, 3rd ed., Graduate Texts in Mathematics, vol. 249, Springer, New York, 2014. ↑4
- [20] R. Hartshorne, *Algebraic geometry*, Graduate Texts in Mathematics, vol. 52, Springer, New York–Heidelberg, 1977. ↑17
- [21] R. A. Horn and C. R. Johnson, *Matrix analysis*, 2nd ed., Cambridge University Press, Cambridge, 2013. ↑12
- [22] L. Lanzani and E. M. Stein, *A note on div curl inequalities*, Math. Res. Lett. **12** (2005), no. 1, 57–61, doi:10.4310/MRL.2005.v12.n1.a6. ↑2
- [23] J.-L. Lions, *Théorèmes de trace et d'interpolation*. I, Ann. Scuola Norm. Sup. Pisa (3) **13** (1959), 389–403; II, Ann. Scuola Norm. Sup. Pisa (3) **14** (1960), 317–331; III, J. Math. Pures Appl. (9) **42** (1963), 195–203; IV, Math. Ann. **151** (1963), 42–56; V, An. Acad. Brasil. Ci. **35** (1963), 1–10. ↑2
- [24] J.-L. Lions and J. Peetre, *Propriétés d'espaces d'interpolation*, C. R. Acad. Sci. Paris **253** (1961), 1747–1749. ↑2
- [25] ———, *Sur une classe d'espaces d'interpolation*, Inst. Hautes Études Sci. Publ. Math. **19** (1964), 5–68. ↑2, 5

- [26] L. H. Loomis and H. Whitney, *An inequality related to the isoperimetric inequality*, Bull. Amer. Math. Soc **55** (1949), 961–962, doi:10.1090/S0002-9904-1949-09320-5. ↑4, 9
- [27] G. G. Lorentz, *Some new functional spaces*, Ann. of Math. (2) **51** (1950), 37–55, doi:10.2307/1969496. ↑4, 5
- [28] V. G. Maz'ya, *Classes of domains and imbedding theorems for function spaces*, Soviet Math. Dokl. **1** (1960), 882–885. ↑
- [29] L. Nirenberg, *On elliptic partial differential equations*, Ann. Scuola Norm. Sup. Pisa (3) **13** (1959), 115–162. ↑1, 8
- [30] R. O'Neil, *Convolution operators and $L(p, q)$ spaces*, Duke Math. J. **30** (1963), 129–142. ↑1
- [31] J. Peetre, *Espaces d'interpolation et théorème de Soboleff*, Ann. Inst. Fourier (Grenoble) **16** (1966), no. fasc. 1, 279–317. ↑1
- [32] S. Poornima, *An embedding theorem for the Sobolev space $W^{1,1}$* , Bull. Sci. Math. (2) **107** (1983), no. 3, 253–259. ↑1
- [33] M. J. Strauss, *Variations of Korn's and Sobolev's equalities*, Partial differential equations (Univ. California, Berkeley, Calif., 1971), Proc. Sympos. Pure Math., vol. XXIII, Amer. Math. Soc., Providence, R.I., 1973, pp. 207–214. ↑2, 15
- [34] L. Tartar, *Imbedding theorems of Sobolev spaces into Lorentz spaces*, Boll. Unione Mat. Ital. Sez. B Artic. Ric. Mat. (8) **1** (1998), no. 3, 479–500. ↑1
- [35] ———, *An introduction to Sobolev spaces and interpolation spaces*, Lecture Notes of the Unione Matematica Italiana, vol. 3, Springer, Berlin; UMI, Bologna, 2007. ↑2, 5
- [36] R. Temam and G. Strang, *Functions of bounded deformation*, Arch. Rational Mech. Anal. **75** (1980/81), no. 1, 7–21. ↑2
- [37] H. Triebel, *Interpolation theory, function spaces, differential operators*, VEB Deutscher Verlag der Wissenschaften, Berlin, 1978. ↑2, 5
- [38] J. Van Schaftingen, *Estimates for L^1 -vector fields*, C. R. Math. Acad. Sci. Paris **339** (2004), no. 3, 181–186, doi:10.1016/j.crma.2004.05.013. ↑2
- [39] ———, *Estimates for L^1 vector fields with a second order condition*, Acad. Roy. Belg. Bull. Cl. Sci. (6) **15** (2004), no. 1-6, 103–112. ↑
- [40] ———, *Limiting fractional and Lorentz space estimates of differential forms*, Proc. Amer. Math. Soc. **138** (2010), no. 1, 235–240, doi:10.1090/S0002-9939-09-10005-9. ↑2, 3
- [41] ———, *Limiting Sobolev inequalities for vector fields and canceling linear differential operators*, J. Eur. Math. Soc. (JEMS) **15** (2013), no. 3, 877–921, doi:10.4171/JEMS/380. ↑2, 3, 4, 15, 18
- [42] ———, *Limiting Bourgain-Brezis estimates for systems of linear differential equations: theme and variations*, J. Fixed Point Theory Appl. **15** (2014), no. 2, 273–297, doi:10.1007/s11784-014-0177-0. ↑2
- [43] W. P. Ziemer, *Weakly differentiable functions: Sobolev spaces and functions of bounded variation*, Graduate Texts in Mathematics, vol. 120, Springer, New York, 1989. ↑5